

Long-time Dynamics and Optimal Control of Diffuse Interface Models for Tumor Growth

E. Rocca

Università degli Studi di Pavia

January 20-26, 2019, Oberwolfach

Workshop on Surface, Bulk, and Geometric Partial Differential Equations:
Interfacial, stochastic, non-local and discrete structures

joint works with Cecilia Cavaterra (Milano), Hao Wu (Fudan),
and Alain Miranville (Poitiers)-Giulio Schimperna (Pavia)



Fondazione Cariplo and Regione Lombardia Grant MEGAsTaR 2016-2019

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Setting

Tumors grown *in vitro* often exhibit “layered” structures:




Figure: Zhang et al. Integr. Biol., 2012, 4, 1072–1080. Scale bar $100\mu\text{m} = 0.1\text{mm}$

Setting

Tumors grown *in vitro* often exhibit “layered” structures:




Figure: Zhang et al. Integr. Biol., 2012, 4, 1072–1080. Scale bar $100\mu\text{m} = 0:1\text{mm}$

A continuum model is introduced with the ansatz:

- sharp interfaces are replaced by narrow transition layers arising due to adhesive forces among the cell species: a **diffuse interface** separates tumor and healthy cell regions
- **proliferating** tumor cells surrounded by (healthy) **host cells**, and a **nutrient** (e.g. glucose).

Setting

Tumors grown *in vitro* often exhibit “layered” structures:





Figure: Zhang et al. Integr. Biol., 2012, 4, 1072–1080. Scale bar $100\mu\text{m} = 0:1\text{mm}$

A continuum model is introduced with the ansatz:

- sharp interfaces are replaced by narrow transition layers arising due to adhesive forces among the cell species: a **diffuse interface** separates tumor and healthy cell regions
- **proliferating** tumor cells surrounded by (healthy) **host cells**, and a **nutrient** (e.g. glucose).

We investigate the long-time dynamics and optimal control problem of a **two-phase** diffuse interface model that describes the growth of a **tumor** in presence of a nutrient and surrounded by **host tissues**.

Diffuse interfaces



The sharp interface S replaced by a
(thickness ϵ) thin transition layer

$\varphi \equiv -1$ in the Healthy tissue phase

$\varphi \equiv 1$ in the Tumor phase

Advantages of diffuse interfaces in tumor growth models

Advantages of diffuse interfaces in tumor growth models

- It eliminates the need to enforce complicated boundary conditions across the tumor/host tissue and other species/species interfaces
- It eliminates the need to explicitly track the position of interfaces, as is required in the sharp interface framework
- The mathematical description remains valid even when the tumor undergoes topological changes (e.g. metastasis)

Advantages of diffuse interfaces in tumor growth models

- It eliminates the need to enforce complicated boundary conditions across the tumor/host tissue and other species/species interfaces
- It eliminates the need to explicitly track the position of interfaces, as is required in the sharp interface framework
- The mathematical description remains valid even when the tumor undergoes topological changes (e.g. metastasis)

Regarding **modeling** of diffuse interface tumor growth we can quote, e.g.,

- Ciarletta, Cristini, Frieboes, Garcke, Hawkins, Hilhorst, Lam, Lowengrub, Oden, Wise, also for their numerical simulations → complex changes in tumor morphologies due to the interactions with nutrients or toxic agents and also due to mechanical stresses

Advantages of diffuse interfaces in tumor growth models

- It eliminates the need to enforce complicated boundary conditions across the tumor/host tissue and other species/species interfaces
- It eliminates the need to explicitly track the position of interfaces, as is required in the sharp interface framework
- The mathematical description remains valid even when the tumor undergoes topological changes (e.g. metastasis)

Regarding **modeling** of diffuse interface tumor growth we can quote, e.g.,

- Ciarletta, Cristini, Frieboes, Garcke, Hawkins, Hilhorst, Lam, Lowengrub, Oden, Wise, also for their numerical simulations → complex changes in tumor morphologies due to the interactions with nutrients or toxic agents and also due to mechanical stresses
- Frieboes, Jin, Chuang, Wise, Lowengrub, Cristini, Garcke, Lam, Nürnberg, Sitka, for the interaction of multiple tumor cell species described by *multiphase mixture models*

Theoretical analysis: two-phase models

- In terms of the theoretical analysis most of the recent literature is restricted to the **two-phase variant**, i.e., to models that only account for the evolution of a tumor surrounded by healthy tissue.

Theoretical analysis: two-phase models

- In terms of the theoretical analysis most of the recent literature is restricted to the **two-phase variant**, i.e., to models that only account for the evolution of a tumor surrounded by healthy tissue.
- In this setting, there is no differentiation among the tumor cells that exhibit heterogeneous growth behavior. Hence this kind of two-phase models are just able to describe the growth of a **young tumor** before the onset of quiescence and necrosis.

Theoretical analysis: two-phase models

- In terms of the theoretical analysis most of the recent literature is restricted to the **two-phase variant**, i.e., to models that only account for the evolution of a tumor surrounded by healthy tissue.
- In this setting, there is no differentiation among the tumor cells that exhibit heterogeneous growth behavior. Hence this kind of two-phase models are just able to describe the growth of a **young tumor** before the onset of quiescence and necrosis.
- Analytical results related to well-posedness, asymptotic limits, but also **optimal control and long-time behavior of solution**, have been established in a number of papers of a number of authors which include: Agosti, Ciarletta, Colli, Frigeri, Garcke, Gilardi, Grasselli, Hilhorst, Lam, Marinoschi, Melchionna, E.R., Scala, Sprekels, Wu, etc...

Theoretical analysis: two-phase models

- In terms of the theoretical analysis most of the recent literature is restricted to the **two-phase variant**, i.e., to models that only account for the evolution of a tumor surrounded by healthy tissue.
- In this setting, there is no differentiation among the tumor cells that exhibit heterogeneous growth behavior. Hence this kind of two-phase models are just able to describe the growth of a **young tumor** before the onset of quiescence and necrosis.
- Analytical results related to well-posedness, asymptotic limits, but also **optimal control and long-time behavior of solution**, have been established in a number of papers of a number of authors which include: Agosti, Ciarletta, Colli, Frigeri, Garcke, Gilardi, Grasselli, Hilhorst, Lam, Marinoschi, Melchionna, E.R., Scala, Sprekels, Wu, etc...
 - ▶ for tumor growth models based on the coupling of Cahn–Hilliard (for the tumor density) and reaction–diffusion (for the nutrient) equations, and
 - ▶ for models of Cahn-Hilliard-Darcy or Cahn-Hilliard-Brinkman type.

Theoretical analysis: two-phase models

- In terms of the theoretical analysis most of the recent literature is restricted to the **two-phase variant**, i.e., to models that only account for the evolution of a tumor surrounded by healthy tissue.
- In this setting, there is no differentiation among the tumor cells that exhibit heterogeneous growth behavior. Hence this kind of two-phase models are just able to describe the growth of a **young tumor** before the onset of quiescence and necrosis.
- Analytical results related to well-posedness, asymptotic limits, but also **optimal control and long-time behavior of solution**, have been established in a number of papers of a number of authors which include: Agosti, Ciarletta, Colli, Frigeri, Garcke, Gilardi, Grasselli, Hilhorst, Lam, Marinoschi, Melchionna, E.R., Scala, Sprekels, Wu, etc...
 - ▶ for tumor growth models based on the coupling of Cahn–Hilliard (for the tumor density) and reaction–diffusion (for the nutrient) equations, and
 - ▶ for models of Cahn-Hilliard-Darcy or Cahn-Hilliard-Brinkman type.

In this talk we concentrate on two recent results on **optimal control** and **long-time behavior of solution**.

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Long-time dynamics and optimal control of a diffuse interface model for tumor growth: joint work with C. Cavaterra and H. Wu (in preparation, 2019)

Long-time dynamics and optimal control of a diffuse interface model for tumor growth: joint work with C. Cavaterra and H. Wu (in preparation, 2019)

- The state system consists of a Cahn-Hilliard type equation for the **tumor cell** fraction and a reaction-diffusion equation for the **nutrient**
 - The possible medication that serves to eliminate tumor cells is in terms of **drugs** and is introduced into the system through the nutrient
 - In this setting, the **control variable** acts as an external source in the nutrient equation
- 1 First, we consider the problem of “**long-time treatment**” under a suitable given source and prove the convergence of any global solution to a single equilibrium as $t \rightarrow +\infty$.
 - 2 Then we consider the “**finite-time treatment**” of tumor, which corresponds to an optimal control problem. Here we also allow the objective cost functional to depend on a free time variable, which represents the **unknown treatment time to be optimized**. We prove the existence of an optimal control and obtain first order necessary optimality conditions for both the drug concentration and the treatment time.

The main modelling idea

One of the main aim of the **control problem is to realize in the best possible way a desired final distribution of the tumor cell**, which is expressed by the target function ϕ_Ω

The main modelling idea

One of the main aim of the **control problem is to realize in the best possible way a desired final distribution of the tumor cell**, which is expressed by the target function ϕ_Ω

By establishing the Lyapunov stability of certain equilibria of the state system (without external source), we see that ϕ_Ω can be taken as a stable configuration, so that the tumor will not grow again once the finite-time treatment is completed

The state system: Cahn–Hilliard + nutrient model with source terms

The state system: Cahn–Hilliard + nutrient model with source terms

The PDE system is an approximation of the model proposed in [HZO: A. Hawkins-Daarud, K.-G. van der Zee and J.-T. Oden (2011)] in $Q := \Omega \times (0, T)$:

$$\begin{aligned}\phi_t - \Delta\mu &= P(\phi)(\sigma - \mu), & \mu &= -\Delta\phi + F'(\phi) \\ \sigma_t - \Delta\sigma &= -P(\phi)(\sigma - \mu) + u\end{aligned}$$

subject to initial and boundary conditions

$$\phi|_{t=0} = \phi_0, \quad \sigma|_{t=0} = \sigma_0, \quad \text{in } \Omega, \quad \partial_\nu\phi = \partial_\nu\mu = \partial_\nu\sigma = 0, \quad \text{on } \partial\Omega \times (0, T)$$

The state system: Cahn–Hilliard + nutrient model with source terms

The PDE system is an approximation of the model proposed in [HZO: A. Hawkins-Daarud, K.-G. van der Zee and J.-T. Oden (2011)] in $Q := \Omega \times (0, T)$:

$$\begin{aligned}\phi_t - \Delta\mu &= P(\phi)(\sigma - \mu), & \mu &= -\Delta\phi + F'(\phi) \\ \sigma_t - \Delta\sigma &= -P(\phi)(\sigma - \mu) + u\end{aligned}$$

subject to initial and boundary conditions

$$\phi|_{t=0} = \phi_0, \quad \sigma|_{t=0} = \sigma_0, \quad \text{in } \Omega, \quad \partial_\nu\phi = \partial_\nu\mu = \partial_\nu\sigma = 0, \quad \text{on } \partial\Omega \times (0, T)$$

- The state variables are:
 - ▶ the tumor cell fraction ϕ : $\phi \simeq 1$ (tumorous phase), $\phi \simeq -1$ (healthy tissue phase)
 - ▶ the nutrient concentration σ : $\sigma \simeq 1$ and $\sigma \simeq 0$ indicate a nutrient-rich or nutrient-poor extracellular water phase
- F is typically a double-well potential with equal minima at $\phi = \pm 1$
- $P \geq 0$ denotes a suitable regular proliferation function
- The choice of **reactive terms** is motivated by the linear phenomenological constitutive laws for chemical reactions
- The **control variable** u serves as an external source in the equation for σ and can be interpreted as a medication

Energy identity

The system turns out to be thermodynamically consistent. In particular, when $u = 0$ the unknown pair (ϕ, σ) is a dissipative gradient flow for the total free energy:

$$\mathcal{E}(\phi, \sigma) = \int_{\Omega} \left[\frac{1}{2} |\nabla \phi|^2 + F(\phi) \right] dx + \frac{1}{2} \int_{\Omega} \sigma^2 dx.$$

Moreover generally, under the presence of the external source u , we observe that any smooth solution (ϕ, σ) to the problem satisfies the following **energy identity**:

$$\frac{d}{dt} \mathcal{E}(\phi, \sigma) + \int_{\Omega} \left[|\nabla \mu|^2 + |\nabla \sigma|^2 + P(\phi)(\mu - \sigma)^2 \right] dx = \int_{\Omega} u \sigma dx,$$

which motivates the twofold aim of the present contribution.

Our results

1. We prove that any global weak solution will converge to a single equilibrium as $t \rightarrow +\infty$ and provide an estimate on the convergence rate.

Our results

1. We prove that any global weak solution will converge to a single equilibrium as $t \rightarrow +\infty$ and provide an estimate on the convergence rate. Our result indicates that after certain medication (or even without medication, i.e., $u = 0$), the tumor will eventually grow to a steady state as time evolves.

Our results

1. We prove that any global weak solution will converge to a single equilibrium as $t \rightarrow +\infty$ and provide an estimate on the convergence rate. Our result indicates that after certain medication (or even without medication, i.e., $u = 0$), **the tumor will eventually grow to a steady state as time evolves**. However, since the potential function F is nonconvex (double-well), the problem may admit **infinite many steady states** so that for the moment one cannot identify which exactly the unique asymptotic limit as $t \rightarrow +\infty$ will be.

Our results

1. We prove that any global weak solution will converge to a single equilibrium as $t \rightarrow +\infty$ and provide an estimate on the convergence rate. Our result indicates that after certain medication (or even without medication, i.e., $u = 0$), **the tumor will eventually grow to a steady state as time evolves**. However, since the potential function F is nonconvex (double-well), the problem may admit **infinite many steady states** so that for the moment one cannot identify which exactly the unique asymptotic limit as $t \rightarrow +\infty$ will be.
2. Denoting by $T \in (0, +\infty)$ a fixed maximal time in which the patient is allowed to undergo a medical treatment, we derive necessary optimality conditions for

(CP) *Minimize the cost functional*

$$\begin{aligned} \mathcal{J}(\phi, \sigma, u, \tau) = & \frac{\beta_Q}{2} \int_0^\tau \int_\Omega |\phi - \phi_Q|^2 \, dx \, dt + \frac{\beta_\Omega}{2} \int_\Omega |\phi(\tau) - \phi_\Omega|^2 \, dx \\ & + \frac{\alpha_Q}{2} \int_0^\tau \int_\Omega |\sigma - \sigma_Q|^2 \, dx \, dt + \frac{\beta_S}{2} \int_\Omega (1 + \phi(\tau)) \, dx + \frac{\beta_u}{2} \int_0^\tau \int_\Omega |u|^2 \, dx \, dt + \beta_T \tau \end{aligned}$$

subject to the state system and the the control constraint

$$u \in \mathcal{U}_{\text{ad}} := \{u \in L^\infty(Q) : u_{\min} \leq u \leq u_{\max} \text{ a. e. in } Q\}, \quad \tau \in (0, T)$$

Comments on the cost functional

$$\begin{aligned}\mathcal{J}(\phi, \sigma, u, \tau) = & \frac{\beta_Q}{2} \int_0^\tau \int_\Omega |\phi - \phi_Q|^2 \, dx \, dt + \frac{\beta_\Omega}{2} \int_\Omega |\phi(\tau) - \phi_\Omega|^2 \, dx \\ & + \frac{\alpha_Q}{2} \int_0^\tau \int_\Omega |\sigma - \sigma_Q|^2 \, dx \, dt + \frac{\beta_S}{2} \int_\Omega (1 + \phi(\tau)) \, dx + \frac{\beta_u}{2} \int_0^T \int_\Omega |u|^2 \, dx \, dt + \beta_{TT}\end{aligned}$$

Comments on the cost functional

$$\begin{aligned} \mathcal{J}(\phi, \sigma, u, \tau) = & \frac{\beta_Q}{2} \int_0^\tau \int_\Omega |\phi - \phi_Q|^2 dx dt + \frac{\beta_\Omega}{2} \int_\Omega |\phi(\tau) - \phi_\Omega|^2 dx \\ & + \frac{\alpha_Q}{2} \int_0^\tau \int_\Omega |\sigma - \sigma_Q|^2 dx dt + \frac{\beta_S}{2} \int_\Omega (1 + \phi(\tau)) dx + \frac{\beta_u}{2} \int_0^\tau \int_\Omega |u|^2 dx dt + \beta_T \tau \end{aligned}$$

- $\tau \in (0, T]$ represents the treatment time of one cycle, i.e., the amount of time the drug is applied to the patient before the period of rest, or the treatment time before surgery, ϕ_Q and σ_Q represent a desired evolution for the tumor cells and for the nutrient, ϕ_Ω stands for desired final distribution of tumor cells
- The first three terms of \mathcal{J} are of standard tracking type and the fourth term of \mathcal{J} measures the size of the tumor at the end of the treatment
- The fifth term penalizes large concentrations of the cytotoxic drugs, and the sixth term of \mathcal{J} penalizes long treatment times

The choice of ϕ_Ω

After the treatment, the ideal situation will be either the tumor is ready for surgery or the tumor will be stable for all time without further medication (i.e., $u = 0$). This goal can be realized by making different choices of the target function ϕ_Ω in the above optimal control problem (CP).

- For the former case, one can simply take ϕ_Ω to be a configuration that is suitable for surgery.
- While for the later case, which is of more interest to us, we want to choose ϕ_Ω as a “stable” configuration of the system, so that the tumor does not grow again once the treatment is complete.

For this purpose, we prove that any local minimizer of the total free energy \mathcal{E} is Lyapunov stable provided that $u = 0$.

The choice of ϕ_Ω

After the treatment, the ideal situation will be either the tumor is ready for surgery or the tumor will be stable for all time without further medication (i.e., $u = 0$). This goal can be realized by making different choices of the target function ϕ_Ω in the above optimal control problem (CP).

- For the former case, one can simply take ϕ_Ω to be a configuration that is suitable for surgery.
- While for the later case, which is of more interest to us, we want to choose ϕ_Ω as a “stable” configuration of the system, so that the tumor does not grow again once the treatment is complete.

For this purpose, we prove that any local minimizer of the total free energy \mathcal{E} is Lyapunov stable provided that $u = 0$. As a consequence, these local energy minimizers serve as possible candidates for the target function ϕ_Ω . Then after completing a successful medication, the tumor will remain close to the chosen stable configuration for all time.

The mathematical difficulties

The mathematical difficulties

The study of long-time behavior is nontrivial, since the nonconvexity of the free energy \mathcal{E} indicates that the set of steady states may have a rather complicated structure.

The mathematical difficulties

The study of long-time behavior is nontrivial, since the nonconvexity of the free energy \mathcal{E} indicates that the set of steady states may have a rather complicated structure.

- For the single Cahn-Hilliard equation this difficulty can be overcome by employing the Łojasiewicz-Simon approach: a key property that plays an important role in the analysis of the Cahn-Hilliard equation is the conservation of mass, i.e.,

$$\int_{\Omega} \phi(t) \, dx = \int_{\Omega} \phi_0 \, dx \quad \text{for } t \geq 0.$$

The mathematical difficulties

The study of long-time behavior is nontrivial, since the nonconvexity of the free energy \mathcal{E} indicates that the set of steady states may have a rather complicated structure.

- For the single Cahn-Hilliard equation this difficulty can be overcome by employing the Łojasiewicz-Simon approach: a key property that plays an important role in the analysis of the Cahn-Hilliard equation is the conservation of mass, i.e.,

$$\int_{\Omega} \phi(t) \, dx = \int_{\Omega} \phi_0 \, dx \quad \text{for } t \geq 0.$$

However, for our coupled system this property no longer holds, which brings us new difficulties in analysis.

The mathematical difficulties

The study of long-time behavior is nontrivial, since the nonconvexity of the free energy \mathcal{E} indicates that the set of steady states may have a rather complicated structure.

- For the single Cahn-Hilliard equation this difficulty can be overcome by employing the Łojasiewicz-Simon approach: a key property that plays an important role in the analysis of the Cahn-Hilliard equation is the conservation of mass, i.e.,

$$\int_{\Omega} \phi(t) \, dx = \int_{\Omega} \phi_0 \, dx \quad \text{for } t \geq 0.$$

However, for our coupled system this property no longer holds, which brings us new difficulties in analysis.

- Besides, quite different from the Cahn-Hilliard-Oono system considered in which the mass $\int_{\Omega} \phi(t) \, dx$ is not preserved due to possible reactions, here in our case it is not obvious how to control the mass changing rate:

$$\frac{d}{dt} \int_{\Omega} \phi \, dx = \int_{\Omega} P(\phi)(\sigma - \mu) \, dx.$$

Similar problem happens to the nutrient as well, that is

$$\frac{d}{dt} \int_{\Omega} \sigma \, dx = - \int_{\Omega} P(\phi)(\sigma - \mu) \, dx + \int_{\Omega} u \, dx.$$

The problem of mass conservation

The problem of mass conservation

- The observation that the **total mass** can be determined by the initial data and the external source:

$$\int_{\Omega} (\phi(t) + \sigma(t)) \, dx = \int_{\Omega} (\phi_0 + \sigma_0) \, dx + \int_0^t \int_{\Omega} u \, dx \, d\tau, \quad \forall t \geq 0$$

allows us to derive a suitable version of the Łojasiewicz-Simon type inequality.

The problem of mass conservation

- The observation that the **total mass** can be determined by the initial data and the external source:

$$\int_{\Omega} (\phi(t) + \sigma(t)) \, dx = \int_{\Omega} (\phi_0 + \sigma_0) \, dx + \int_0^t \int_{\Omega} u \, dx \, d\tau, \quad \forall t \geq 0$$

allows us to derive a suitable version of the Łojasiewicz-Simon type inequality.

- On the other hand, we can control the mass changing rates of ϕ and σ by using the extra dissipation related to reactive terms in the basic energy law, i.e.,

$$\int_{\Omega} P(\phi)(\mu - \sigma)^2 \, dx.$$

The problem of mass conservation

- The observation that the **total mass** can be determined by the initial data and the external source:

$$\int_{\Omega} (\phi(t) + \sigma(t)) \, dx = \int_{\Omega} (\phi_0 + \sigma_0) \, dx + \int_0^t \int_{\Omega} u \, dx \, d\tau, \quad \forall t \geq 0$$

allows us to derive a suitable version of the Łojasiewicz-Simon type inequality.

- On the other hand, we can control the mass changing rates of ϕ and σ by using the extra dissipation related to reactive terms in the basic energy law, i.e.,
$$\int_{\Omega} P(\phi)(\mu - \sigma)^2 \, dx.$$
- Based on the above mentioned special structure of the system, by introducing a new version of Łojasiewicz-Simon inequality we are able to prove that every global weak solution (ϕ, σ) of the problem will converge to a certain single equilibrium $(\phi_{\infty}, \sigma_{\infty})$ as $t \rightarrow +\infty$ and, moreover, we obtain a polynomial decay of the solution.

The problem of mass conservation

- The observation that the **total mass** can be determined by the initial data and the external source:

$$\int_{\Omega} (\phi(t) + \sigma(t)) \, dx = \int_{\Omega} (\phi_0 + \sigma_0) \, dx + \int_0^t \int_{\Omega} u \, dx \, d\tau, \quad \forall t \geq 0$$

allows us to derive a suitable version of the Łojasiewicz-Simon type inequality.

- On the other hand, we can control the mass changing rates of ϕ and σ by using the extra dissipation related to reactive terms in the basic energy law, i.e.,
$$\int_{\Omega} P(\phi)(\mu - \sigma)^2 \, dx.$$
- Based on the above mentioned special structure of the system, by introducing a new version of Łojasiewicz-Simon inequality we are able to prove that every global weak solution (ϕ, σ) of the problem will converge to a certain single equilibrium $(\phi_{\infty}, \sigma_{\infty})$ as $t \rightarrow +\infty$ and, moreover, we obtain a polynomial decay of the solution.
- Besides, a nontrivial application of the Łojasiewicz-Simon approach further leads to the Lyapunov stability of local minimizers of the free energy \mathcal{E} (we only consider the case $u = 0$ for the sake of simplicity).

Comparison with other results in the literature

- To the best of our knowledge, the only contribution in the study of long-time behavior for this problem is given in [FGR: Frigeri, Grasselli, R. (2015)] with $u = 0$, where, however, the main focus is the **existence of a global attractor**

Comparison with other results in the literature

- To the best of our knowledge, the only contribution in the study of long-time behavior for this problem is given in [FGR: Frigeri, Grasselli, R. (2015)] with $u = 0$, where, however, the main focus is the **existence of a global attractor**
- In the context of PDE constraint optimal control for diffuse interface tumor models, in the literature we have basically two recent works:

Comparison with other results in the literature

- To the best of our knowledge, the only contribution in the study of long-time behavior for this problem is given in [FGR: Frigeri, Grasselli, R. (2015)] with $u = 0$, where, however, the main focus is the **existence of a global attractor**
- In the context of PDE constraint optimal control for diffuse interface tumor models, in the literature we have basically two recent works:
 - 1 [CGRS: Colli, Gilardi, R., Sprekels (2017)] where the objective functional is with the special (simpler) choices $\beta_S = \beta_T = \alpha_Q = 0$, and the state system is exactly the same but **no dependence on τ** is studied.

Comparison with other results in the literature

- To the best of our knowledge, the only contribution in the study of long-time behavior for this problem is given in [FGR: Frigeri, Grasselli, R. (2015)] with $u = 0$, where, however, the main focus is the **existence of a global attractor**
- In the context of PDE constraint optimal control for diffuse interface tumor models, in the literature we have basically two recent works:
 - 1 [CGRS: Colli, Gilardi, R., Sprekels (2017)] where the objective functional is with the special (simpler) choices $\beta_S = \beta_T = \alpha_Q = 0$, and the state system is exactly the same but **no dependence on τ** is studied.
 - 2 [GLR: Garcke, Lam, R. (2017)] where a **different diffuse interface model** is studied. There the distributed control appears in the ϕ equation, which is a Cahn-Hilliard type equation with a source of mass on the right hand side, but not depending on μ . Due to the presence of **the control in the Cahn-Hilliard equation, in [GLR] only the case of a regularized objective cost functional** can be analyzed for bounded controls.

Comparison with other results in the literature

- To the best of our knowledge, the only contribution in the study of long-time behavior for this problem is given in [FGR: Frigeri, Grasselli, R. (2015)] with $u = 0$, where, however, the main focus is the **existence of a global attractor**
- In the context of PDE constraint optimal control for diffuse interface tumor models, in the literature we have basically two recent works:
 - 1 [CGRS: Colli, Gilardi, R., Sprekels (2017)] where the objective functional is with the special (simpler) choices $\beta_S = \beta_T = \alpha_Q = 0$, and the state system is exactly the same but **no dependence on τ** is studied.
 - 2 [GLR: Garcke, Lam, R. (2017)] where a **different diffuse interface model** is studied. There the distributed control appears in the ϕ equation, which is a Cahn-Hilliard type equation with a source of mass on the right hand side, but not depending on μ . Due to the presence of **the control in the Cahn-Hilliard equation, in [GLR] only the case of a regularized objective cost functional** can be analyzed for bounded controls.

With our work we aim to provide a contribution to the theory of free terminal time optimal control in the context of diffuse interface tumor models, where the control is applied in the nutrient equation.

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness**
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Well-posedness (cf, [CGRS, Theorem 2.1])

Well-posedness (cf, [CGRS, Theorem 2.1])

Let $\phi_0 \in H_N^2(\Omega) \cap H^3(\Omega)$ and $\sigma_0 \in H^1(\Omega)$ and assume that

- (P1) $P \in C^2(\mathbb{R})$ is nonnegative. There exist $\alpha_1 > 0$ and some $q \in [1, 4]$ such that, for all $s \in \mathbb{R}$, $|P'(s)| \leq \alpha_1(1 + |s|^{q-1})$
- (F1) $F = F_0 + F_1$, with $F_0, F_1 \in C^5(\mathbb{R})$. There exist $\alpha_i > 0$ and $r \in [2, 6)$ such that $|F_1''(s)| \leq \alpha_2$, $\alpha_3(1 + |s|^{r-2}) \leq F_0''(s) \leq \alpha_4(1 + |s|^{r-2})$, $F(s) \geq \alpha_5|s| - \alpha_6 \quad \forall s \in \mathbb{R}$
- (U1) For any $T > 0$, $u \in L^2(0, T; L^2(\Omega))$.

Well-posedness (cf, [CGRS, Theorem 2.1])

Let $\phi_0 \in H_N^2(\Omega) \cap H^3(\Omega)$ and $\sigma_0 \in H^1(\Omega)$ and assume that

- (P1) $P \in C^2(\mathbb{R})$ is nonnegative. There exist $\alpha_1 > 0$ and some $q \in [1, 4]$ such that, for all $s \in \mathbb{R}$, $|P'(s)| \leq \alpha_1(1 + |s|^{q-1})$
- (F1) $F = F_0 + F_1$, with $F_0, F_1 \in C^5(\mathbb{R})$. There exist $\alpha_i > 0$ and $r \in [2, 6)$ such that $|F_1''(s)| \leq \alpha_2$, $\alpha_3(1 + |s|^{r-2}) \leq F_0''(s) \leq \alpha_4(1 + |s|^{r-2})$, $F(s) \geq \alpha_5|s| - \alpha_6 \quad \forall s \in \mathbb{R}$
- (U1) For any $T > 0$, $u \in L^2(0, T; L^2(\Omega))$. Then

Theorem (Strong solutions)

(1) For every $T > 0$, the state system admits a unique strong solution:

$$\begin{aligned} & \|\phi\|_{L^\infty(0, T; H^3(\Omega)) \cap L^2(0, T; H^4(\Omega)) \cap H^1(0, T; H^1(\Omega))} + \|\mu\|_{L^\infty(0, T; H^1(\Omega)) \cap L^2(0, T; H^2(\Omega))} \\ & + \|\sigma\|_{C([0, T]; H^1(\Omega)) \cap L^2(0, T; H_N^2(\Omega)) \cap H^1(0, T; L^2(\Omega))} \leq K_1. \end{aligned}$$

Well-posedness (cf, [CGRS, Theorem 2.1])

Let $\phi_0 \in H_N^2(\Omega) \cap H^3(\Omega)$ and $\sigma_0 \in H^1(\Omega)$ and assume that

- (P1) $P \in C^2(\mathbb{R})$ is nonnegative. There exist $\alpha_1 > 0$ and some $q \in [1, 4]$ such that, for all $s \in \mathbb{R}$, $|P'(s)| \leq \alpha_1(1 + |s|^{q-1})$
- (F1) $F = F_0 + F_1$, with $F_0, F_1 \in C^5(\mathbb{R})$. There exist $\alpha_i > 0$ and $r \in [2, 6)$ such that
- $$|F_1''(s)| \leq \alpha_2, \quad \alpha_3(1 + |s|^{r-2}) \leq F_0''(s) \leq \alpha_4(1 + |s|^{r-2}), \quad F(s) \geq \alpha_5|s| - \alpha_6 \quad \forall s \in \mathbb{R}$$
- (U1) For any $T > 0$, $u \in L^2(0, T; L^2(\Omega))$. Then

Theorem (Strong solutions)

(1) For every $T > 0$, the state system admits a unique strong solution:

$$\begin{aligned} & \|\phi\|_{L^\infty(0, T; H^3(\Omega)) \cap L^2(0, T; H^4(\Omega)) \cap H^1(0, T; H^1(\Omega))} + \|\mu\|_{L^\infty(0, T; H^1(\Omega)) \cap L^2(0, T; H^2(\Omega))} \\ & + \|\sigma\|_{C([0, T]; H^1(\Omega)) \cap L^2(0, T; H_N^2(\Omega)) \cap H^1(0, T; L^2(\Omega))} \leq K_1. \end{aligned}$$

(2) Let (ϕ_i, σ_i) be two strong solutions. Then there exists a constant $K_2 > 0$, depending on $\|u_i\|_{L^2(0, T; L^2)}$, Ω , T , $\|\phi_0\|_{H^3}$ and $\|\sigma_0\|_{H^1}$, such that

$$\begin{aligned} & \|\phi_1 - \phi_2\|_{L^\infty(0, T; H^1) \cap L^2(0, T; H^3) \cap H^1(0, T; (H^1)')} + \|\mu_1 - \mu_2\|_{L^2(0, T; H^1)} \\ & + \|\sigma_1 - \sigma_2\|_{C([0, T]; H^1) \cap L^2(0, T; H^2) \cap H^1(0, T; L^2)} \leq K_2 \|\mathbf{u}_1 - \mathbf{u}_2\|_{L^2(0, T; L^2)}. \end{aligned}$$

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics**
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Long-term dynamics

We make the following additional assumptions:

(P2) $P(s) > 0$, for all $s \in \mathbb{R}$

(F2) $F(s)$ is real analytic, for all $s \in \mathbb{R}$

(U2) $u \in L^1(0, +\infty; L^2(\Omega)) \cap L^2(0, +\infty; L^2(\Omega))$ and satisfies the decay condition

$$\sup_{t \geq 0} (1+t)^{3+\rho} \|u(t)\|_{L^2(\Omega)} < +\infty, \quad \text{for some } \rho > 0.$$

Long-term dynamics

We make the following additional assumptions:

- (P2) $P(s) > 0$, for all $s \in \mathbb{R}$
- (F2) $F(s)$ is real analytic, for all $s \in \mathbb{R}$
- (U2) $u \in L^1(0, +\infty; L^2(\Omega)) \cap L^2(0, +\infty; L^2(\Omega))$ and satisfies the decay condition

$$\sup_{t \geq 0} (1+t)^{3+\rho} \|u(t)\|_{L^2(\Omega)} < +\infty, \quad \text{for some } \rho > 0.$$

Theorem (1. The stationary problem)

For any $\phi_0 \in H^1(\Omega)$, $\sigma_0 \in L^2(\Omega)$, the state system admits a unique global weak solution (ϕ, μ, σ) : $\lim_{t \rightarrow +\infty} (\|\phi(t) - \phi_\infty\|_{H^2(\Omega)} + \|\sigma(t) - \sigma_\infty\|_{L^2(\Omega)} + \|\mu(t) - \mu_\infty\|_{L^2(\Omega)}) = 0$, where $(\phi_\infty, \mu_\infty, \sigma_\infty)$ satisfies the stationary problem

$$\left\{ \begin{array}{ll} -\Delta \phi_\infty + F'(\phi_\infty) = \mu_\infty, & \text{in } \Omega \\ \partial_\nu \phi_\infty = 0, & \text{on } \partial\Omega \\ \int_\Omega (\phi_\infty + \sigma_\infty) dx = \int_\Omega (\phi_0 + \sigma_0) dx + \int_0^{+\infty} \int_\Omega u dx dt \end{array} \right.$$

with μ_∞ and σ_∞ being two constants given by $\sigma_\infty = \mu_\infty = |\Omega|^{-1} \int_\Omega F'(\phi_\infty) dx$.

The convergence rate

Theorem (2. Convergence rate)

Moreover, under the same assumptions, the following estimates on convergence rate hold

$$\|\phi(t) - \phi_\infty\|_{H^1(\Omega)} + \|\sigma(t) - \sigma_\infty\|_{L^2(\Omega)} \leq C(1+t)^{-\min\{\frac{\theta}{1-2\theta}, \frac{\rho}{2}\}}, \quad \forall t \geq 0,$$

$$\|\mu(t) - \mu_\infty\|_{L^2(\Omega)} \leq C(1+t)^{-\frac{1}{2} \min\{\frac{\theta}{1-2\theta}, \frac{\rho}{2}\}}, \quad \forall t \geq 0,$$

where $C > 0$ is a constant depending on $\|\phi_0\|_{H^1(\Omega)}$, $\|\sigma_0\|_{L^2(\Omega)}$, $\|\phi_\infty\|_{H^1(\Omega)}$, $\|u\|_{L^1(0,+\infty;L^2(\Omega))}$, $\|u\|_{L^2(0,+\infty;L^2(\Omega))}$ and Ω ; $\theta \in (0, \frac{1}{2})$ is a constant depending on ϕ_∞ .

An idea of the proof

An idea of the proof

The proof consists of several steps:

- We first derive some uniform-in-time a priori estimates on the solution (ϕ, μ, σ)

An idea of the proof

The proof consists of several steps:

- We first derive some uniform-in-time a priori estimates on the solution (ϕ, μ, σ)
- Then we give a characterization on the ω -limit

$$\omega(\phi_0, \sigma_0) = \{(\phi_\infty, \sigma_\infty) \in (H_N^2(\Omega) \cap H^3(\Omega)) \times H^1(\Omega) : \exists \{t_n\} \nearrow +\infty \text{ such that } (\phi(t_n), \sigma(t_n)) \rightarrow (\phi_\infty, \sigma_\infty) \text{ in } H^2(\Omega) \times L^2(\Omega)\}.$$

And we have the following result

Theorem (3. The ω -limit)

Assume (P1), (F1), (U2). For any initial datum $(\phi_0, \sigma_0) \in H^1(\Omega) \times L^2(\Omega)$, the associated ω -limit set $\omega(\phi_0, \sigma_0)$ is non-empty. For any element $(\phi_\infty, \sigma_\infty) \in \omega(\phi_0, \sigma_0)$, σ_∞ is a constant and $(\phi_\infty, \sigma_\infty)$ satisfies the stationary problem. Besides, μ_∞ is a constant given by $|\Omega|^{-1} \int_\Omega F'(\phi_\infty) dx$ and the following relation holds

$$P(\phi_\infty)(\sigma_\infty - \mu_\infty) = 0, \quad \text{a.e. in } \Omega.$$

And the positivity of P entails immediately also $\sigma_\infty = \mu_\infty$.

- Finally, we prove the convergence of the trajectories and polynomial decay by means of a proper Łojasiewicz–Simon inequality:

- Finally, we prove the convergence of the trajectories and polynomial decay by means of a proper Łojasiewicz–Simon inequality: Given any initial datum $(\phi_0, \sigma_0) \in H^1(\Omega) \times L^2(\Omega)$ and source term u satisfying (U2), we denote by

$$m_\infty := |\Omega|^{-1} \left(\int_{\Omega} (\phi_0 + \sigma_0) dx + \int_0^{+\infty} \int_{\Omega} u dx dt \right)$$

the total mass at infinity time. Then we are able to derive the following

Theorem (Łojasiewicz–Simon Inequality)

Let (F1), (F2), (P1), (P2) and (U2) be satisfied. Suppose that $(\phi_\infty, \mu_\infty, \sigma_\infty)$ is a solution to the elliptic stationary problem. Then there exist constants $\theta \in (0, \frac{1}{2})$ and $\beta > 0$, depending on ϕ_∞ , m_∞ and Ω , such that for any $(\phi, \sigma) \in H_N^2(\Omega) \times H^1(\Omega)$ satisfying

$$\|\phi - \phi_\infty\|_{H^1(\Omega)} < \beta,$$

$$\int_{\Omega} (\phi + \sigma) dx + m_u |\Omega| = \int_{\Omega} (\phi_\infty + \sigma_\infty) dx = m_\infty |\Omega|,$$

where m_u is a certain constant fulfilling $|m_u| \leq |\Omega|^{-\frac{1}{2}} \|u\|_{L^1(0,+\infty;L^2(\Omega))}$, then we have

$$\begin{aligned} \|\mu - \bar{\mu}\|_{(H^1(\Omega))'} + C \|\nabla \sigma\|_{L^2(\Omega)} + C \|\sqrt{P(\phi)}(\mu - \sigma)\|_{L^2(\Omega)} + C |m_u|^{\frac{1}{2}} \\ \geq |\mathcal{E}(\phi, \sigma) - \mathcal{E}(\phi_\infty, \sigma_\infty)|^{1-\theta}, \quad \text{where} \end{aligned}$$

$\mu = -\Delta \phi + F'(\phi)$ and $C > 0$ depends on Ω , ϕ_∞ , m_∞ , $\|\phi\|_{H^2(\Omega)}$, $\|\sigma\|_{H^1(\Omega)}$, $\|u\|_{L^1(0,+\infty;L^2(\Omega))}$.

Energy minimizers with $u = 0$

Energy minimizers with $u = 0$

Let us now **assume** $u = 0$. Then it follows that the total mass of the system is now conserved:

$$\int_{\Omega} (\phi(t) + \sigma(t)) \, dx = \int_{\Omega} (\phi_0 + \sigma_0) \, dx, \quad \forall t \geq 0.$$

Energy minimizers with $u = 0$

Let us now **assume** $u = 0$. Then it follows that the total mass of the system is now conserved:

$$\int_{\Omega} (\phi(t) + \sigma(t)) \, dx = \int_{\Omega} (\phi_0 + \sigma_0) \, dx, \quad \forall t \geq 0.$$

Let $m \in \mathbb{R}$ be an arbitrary given constant. Set

$$\mathcal{Z}_m = \left\{ (\phi, \sigma) \in H^1(\Omega) \times L^2(\Omega) : \int_{\Omega} (\phi + \sigma) \, dx = |\Omega| m \right\}.$$

Energy minimizers with $u = 0$

Let us now **assume** $u = 0$. Then it follows that the total mass of the system is now conserved:

$$\int_{\Omega} (\phi(t) + \sigma(t)) \, dx = \int_{\Omega} (\phi_0 + \sigma_0) \, dx, \quad \forall t \geq 0.$$

Let $m \in \mathbb{R}$ be an arbitrary given constant. Set

$$\mathcal{Z}_m = \left\{ (\phi, \sigma) \in H^1(\Omega) \times L^2(\Omega) : \int_{\Omega} (\phi + \sigma) \, dx = |\Omega|m \right\}.$$

Any $(\phi^*, \sigma^*) \in \mathcal{Z}_m$ is called

- a *local energy minimizer* of the total energy

$$\mathcal{E}(\phi, \sigma) = \int_{\Omega} \left[\frac{1}{2} |\nabla \phi|^2 + F(\phi) \right] \, dx + \frac{1}{2} \int_{\Omega} \sigma^2 \, dx$$

if there exists a constant $\chi > 0$ such that $\mathcal{E}(\phi^*, \sigma^*) \leq \mathcal{E}(\phi, \sigma)$, for all $(\phi, \sigma) \in \mathcal{Z}_m$ satisfying $\|(\phi - \phi^*, \sigma - \sigma^*)\|_{H^1(\Omega) \times L^2(\Omega)} < \chi$

- If $\chi = +\infty$, then (ϕ^*, σ^*) is called a *global energy minimizer* of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m .

We first derive some properties for the critical points of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m .

We first derive some properties for the critical points of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . For any given $m \in \mathbb{R}$, we consider the following stationary problem for (ϕ, μ, σ)

$$\left\{ \begin{array}{ll} -\Delta\phi + F'(\phi) = \mu, & \text{in } \Omega, \\ \partial_\nu\phi = 0, & \text{on } \partial\Omega, \\ \int_\Omega (\phi + \sigma) \, dx = |\Omega|m, & \end{array} \right.$$

where μ and σ are constants given by $\sigma = \mu = |\Omega|^{-1} \int_\Omega F'(\phi) \, dx$.

We first derive some properties for the critical points of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . For any given $m \in \mathbb{R}$, we consider the following stationary problem for (ϕ, μ, σ)

$$\begin{cases} -\Delta\phi + F'(\phi) = \mu, & \text{in } \Omega, \\ \partial_\nu\phi = 0, & \text{on } \partial\Omega, \\ \int_\Omega (\phi + \sigma) \, dx = |\Omega|m, \end{cases}$$

where μ and σ are constants given by $\sigma = \mu = |\Omega|^{-1} \int_\Omega F'(\phi) \, dx$.

Theorem (4. Critical points)

Let assumption (F1) be satisfied. Then we have:

We first derive some properties for the critical points of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . For any given $m \in \mathbb{R}$, we consider the following stationary problem for (ϕ, μ, σ)

$$\begin{cases} -\Delta\phi + F'(\phi) = \mu, & \text{in } \Omega, \\ \partial_\nu\phi = 0, & \text{on } \partial\Omega, \\ \int_\Omega (\phi + \sigma) \, dx = |\Omega|m, \end{cases}$$

where μ and σ are constants given by $\sigma = \mu = |\Omega|^{-1} \int_\Omega F'(\phi) \, dx$.

Theorem (4. Critical points)

Let assumption (F1) be satisfied. Then we have:

- (1) If $(\phi^*, \sigma^*) \in H_N^2(\Omega) \times \mathbb{R}$ is a strong solution to the stationary problem above, then (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . Conversely, if (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m , then $\phi^* \in H_N^2(\Omega)$, $\sigma^* \in \mathbb{R}$ satisfy the stationary problem above

We first derive some properties for the critical points of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . For any given $m \in \mathbb{R}$, we consider the following stationary problem for (ϕ, μ, σ)

$$\begin{cases} -\Delta\phi + F'(\phi) = \mu, & \text{in } \Omega, \\ \partial_\nu\phi = 0, & \text{on } \partial\Omega, \\ \int_\Omega (\phi + \sigma) \, dx = |\Omega|m, \end{cases}$$

where μ and σ are constants given by $\sigma = \mu = |\Omega|^{-1} \int_\Omega F'(\phi) \, dx$.

Theorem (4. Critical points)

Let assumption (F1) be satisfied. Then we have:

- (1) If $(\phi^*, \sigma^*) \in H_N^2(\Omega) \times \mathbb{R}$ is a strong solution to the stationary problem above, then (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . Conversely, if (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m , then $\phi^* \in H_N^2(\Omega)$, $\sigma^* \in \mathbb{R}$ satisfy the stationary problem above
- (2) If (ϕ^*, σ^*) is a local energy minimizer of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m , then (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$.

We first derive some properties for the critical points of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . For any given $m \in \mathbb{R}$, we consider the following stationary problem for (ϕ, μ, σ)

$$\begin{cases} -\Delta\phi + F'(\phi) = \mu, & \text{in } \Omega, \\ \partial_\nu\phi = 0, & \text{on } \partial\Omega, \\ \int_\Omega (\phi + \sigma) \, dx = |\Omega|m, \end{cases}$$

where μ and σ are constants given by $\sigma = \mu = |\Omega|^{-1} \int_\Omega F'(\phi) \, dx$.

Theorem (4. Critical points)

Let assumption (F1) be satisfied. Then we have:

- (1) If $(\phi^*, \sigma^*) \in H_N^2(\Omega) \times \mathbb{R}$ is a strong solution to the stationary problem above, then (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m . Conversely, if (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m , then $\phi^* \in H_N^2(\Omega)$, $\sigma^* \in \mathbb{R}$ satisfy the stationary problem above
- (2) If (ϕ^*, σ^*) is a local energy minimizer of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m , then (ϕ^*, σ^*) is a critical point of $\mathcal{E}(\phi, \sigma)$.
- (3) The functional $\mathcal{E}(\phi, \sigma)$ has at least one minimizer $(\phi^*, \sigma^*) \in \mathcal{Z}_m$ such that

$$\mathcal{E}(\phi^*, \sigma^*) = \inf_{(\phi, \sigma) \in \mathcal{Z}_m} \mathcal{E}(\phi, \sigma)$$

Lyapunov Stability with $u = 0$

Lyapunov Stability with $u = 0$

Then, we can get our main result on long-term dynamics:

Lyapunov Stability with $u = 0$

Then, we can get our main result on long-term dynamics:

Theorem (5. Lyapunov stability)

Assume that (F1), (F2), (P1), (P2) are satisfied and $u = 0$. Given $m \in \mathbb{R}$, let (ϕ^*, σ^*) be a local energy minimizer in \mathcal{Z}_m of

$$\mathcal{E}(\phi, \sigma) = \int_{\Omega} \left[\frac{1}{2} |\nabla \phi|^2 + F(\phi) \right] dx + \frac{1}{2} \int_{\Omega} \sigma^2 dx.$$

Lyapunov Stability with $u = 0$

Then, we can get our main result on long-term dynamics:

Theorem (5. Lyapunov stability)

Assume that (F1), (F2), (P1), (P2) are satisfied and $u = 0$. Given $m \in \mathbb{R}$, let (ϕ^*, σ^*) be a local energy minimizer in \mathcal{Z}_m of

$$\mathcal{E}(\phi, \sigma) = \int_{\Omega} \left[\frac{1}{2} |\nabla \phi|^2 + F(\phi) \right] dx + \frac{1}{2} \int_{\Omega} \sigma^2 dx.$$

Then, for any $\epsilon > 0$, there exists a constant $\eta \in (0, 1)$ such that for arbitrary initial datum $(\phi_0, \sigma_0) \in (H_N^2(\Omega) \cap H^3(\Omega)) \times H^1(\Omega)$ satisfying $\int_{\Omega} (\phi_0 + \sigma_0) dx = |\Omega|m$ and $\|\phi_0 - \phi^*\|_{H^1(\Omega)} + \|\sigma_0 - \sigma^*\|_{L^2(\Omega)} \leq \eta$, the state system admits a unique global strong solution (ϕ, σ) such that

$$\|\phi(t) - \phi^*\|_{H^1(\Omega)} + \|\sigma(t) - \sigma^*\|_{L^2(\Omega)} \leq \epsilon, \quad \forall t \geq 0.$$

Namely, any local energy minimizer of $\mathcal{E}(\phi, \sigma)$ in \mathcal{Z}_m is locally Lyapunov stable.

Conclusions on long-term dynamics

- The result on long-time behavior derived in Theorem 1 and 2 can be applied to the global strong solution obtained in Theorem 5
- Although it is still not obvious to identify the asymptotic limit $(\phi_\infty, \sigma_\infty)$, we are able to conclude that $(\phi_\infty, \sigma_\infty)$ also satisfies

$$\|\phi_\infty - \phi^*\|_{H^1(\Omega)} + \|\sigma_\infty - \sigma^*\|_{L^2(\Omega)} \leq \epsilon$$

- In particular, if (ϕ^*, σ^*) is an isolated local energy minimizer then it is locally asymptotic stable

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem**
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Assumptions for the optimal control problem

In this section we study the optimal control problem

(CP) *Minimize the cost functional*

$$\begin{aligned} \mathcal{J}(\phi, \sigma, u, \tau) = & \frac{\beta_Q}{2} \int_0^\tau \int_\Omega |\phi - \phi_Q|^2 \, dx \, dt + \frac{\beta_\Omega}{2} \int_\Omega |\phi(\tau) - \phi_\Omega|^2 \, dx \\ & + \frac{\alpha_Q}{2} \int_0^\tau \int_\Omega |\sigma - \sigma_Q|^2 \, dx \, dt + \frac{\beta_S}{2} \int_\Omega (1 + \phi(\tau)) \, dx + \frac{\beta_u}{2} \int_0^\tau \int_\Omega |u|^2 \, dx \, dt + \beta_T \tau \end{aligned}$$

subject to the state system and the the control constraint

$$u \in \mathcal{U}_{\text{ad}} := \{u \in L^\infty(Q) : u_{\min} \leq u \leq u_{\max} \text{ a. e. in } Q\}, \quad \tau \in (0, T),$$

Assumptions for the optimal control problem

In this section we study the optimal control problem

(CP) *Minimize the cost functional*

$$\begin{aligned} \mathcal{J}(\phi, \sigma, u, \tau) = & \frac{\beta_Q}{2} \int_0^\tau \int_\Omega |\phi - \phi_Q|^2 \, dx \, dt + \frac{\beta_\Omega}{2} \int_\Omega |\phi(\tau) - \phi_\Omega|^2 \, dx \\ & + \frac{\alpha_Q}{2} \int_0^\tau \int_\Omega |\sigma - \sigma_Q|^2 \, dx \, dt + \frac{\beta_S}{2} \int_\Omega (1 + \phi(\tau)) \, dx + \frac{\beta_u}{2} \int_0^T \int_\Omega |u|^2 \, dx \, dt + \beta_T \tau \end{aligned}$$

subject to the state system and the the control constraint

$$u \in \mathcal{U}_{\text{ad}} := \{u \in L^\infty(Q) : u_{\min} \leq u \leq u_{\max} \text{ a. e. in } Q\}, \quad \tau \in (0, T),$$

where $T \in (0, +\infty)$ is a fixed maximal time. We assume:

Assumptions for the optimal control problem

In this section we study the optimal control problem

(CP) *Minimize the cost functional*

$$\begin{aligned} \mathcal{J}(\phi, \sigma, u, \tau) = & \frac{\beta_Q}{2} \int_0^\tau \int_\Omega |\phi - \phi_Q|^2 \, dx \, dt + \frac{\beta_\Omega}{2} \int_\Omega |\phi(\tau) - \phi_\Omega|^2 \, dx \\ & + \frac{\alpha_Q}{2} \int_0^\tau \int_\Omega |\sigma - \sigma_Q|^2 \, dx \, dt + \frac{\beta_S}{2} \int_\Omega (1 + \phi(\tau)) \, dx + \frac{\beta_u}{2} \int_0^\tau \int_\Omega |u|^2 \, dx \, dt + \beta_T \tau \end{aligned}$$

subject to the state system and the the control constraint

$$u \in \mathcal{U}_{\text{ad}} := \{u \in L^\infty(Q) : u_{\min} \leq u \leq u_{\max} \text{ a. e. in } Q\}, \quad \tau \in (0, T),$$

where $T \in (0, +\infty)$ is a fixed maximal time. We assume:

- (C1) $\beta_Q, \beta_\Omega, \beta_S, \beta_u, \beta_T, \alpha_Q$ are nonnegative constants but not all zero.
- (C2) $\phi_Q \in L^2(Q)$, $\phi_\Omega, \sigma_\Omega \in L^2(\Omega)$, $u_{\min}, u_{\max} \in L^\infty(Q)$, and $u_{\min} \leq u_{\max}$, a.e. in Q .
- (C3) Let \mathcal{U}_R be an open set in $L^2(Q)$: $\mathcal{U}_{\text{ad}} \subset \mathcal{U}_R$ and $\|u\|_{L^2(Q)} \leq R$, for all $u \in \mathcal{U}_R$.

Differences with respect to [CGRS: Colli, Gilardi, R., Sprekels (2017)] and [GLR: Garcke, Lam, R. (2017)]

- Here we generalize the problem of [CGRS] by adding the dependence on τ in the cost functional \mathcal{J} .
- Moreover, we can consider the τ -dependent terms in ϕ in the cost functional

$$\begin{aligned}\mathcal{J}(\phi, \sigma, u, \tau) &= \frac{\beta_Q}{2} \int_0^\tau \int_\Omega |\phi - \phi_Q|^2 \, dx \, dt + \frac{\beta_\Omega}{2} \int_\Omega |\phi(\tau) - \phi_\Omega|^2 \, dx \\ &+ \frac{\alpha_Q}{2} \int_0^\tau \int_\Omega |\sigma - \sigma_Q|^2 \, dx \, dt + \frac{\beta_S}{2} \int_\Omega (1 + \phi(\tau)) \, dx + \frac{\beta_u}{2} \int_0^T \int_\Omega |u|^2 \, dx \, dt + \beta_{T\tau}\end{aligned}$$

which we were not able to handle in the paper [GLR], mainly due to the fact that the control u here is in the nutrient equation instead of in the phase equation. Due to this fact, we can enhance the regularity results on ϕ without affecting the regularity of the control u .

Existence of an optimal control

Existence of an optimal control

From the well-posedness results it follows that the *control-to-state operator* \mathcal{S}

$$u \mapsto \mathcal{S}(u) := (\phi, \mu, \sigma)$$

is well-defined and Lipschitz continuous as a mapping from $\mathcal{U}_R \subset L^2(Q)$ into the following space

$$(L^\infty(0, T; (H^1(\Omega))') \cap L^2(0, T; H^1(\Omega))) \times L^2(0, T; (H^1(\Omega))') \times (L^\infty(0, T; (H^1(\Omega))') \cap L^2(Q)).$$

Existence of an optimal control

From the well-posedness results it follows that the *control-to-state operator* \mathcal{S}

$$u \mapsto \mathcal{S}(u) := (\phi, \mu, \sigma)$$

is well-defined and Lipschitz continuous as a mapping from $\mathcal{U}_R \subset L^2(Q)$ into the following space

$$(L^\infty(0, T; (H^1(\Omega))') \cap L^2(0, T; H^1(\Omega))) \times L^2(0, T; (H^1(\Omega))') \times (L^\infty(0, T; (H^1(\Omega))') \cap L^2(Q)).$$

The triplet (ϕ, μ, σ) is the unique weak solution to the state system with data (ϕ_0, σ_0, u) over the time interval $[0, T]$. For convenience, we use the notations $\phi = \mathcal{S}_1(u)$ and $\sigma = \mathcal{S}_3(u)$ for the first and third component of $\mathcal{S}(u)$.

Existence of an optimal control

From the well-posedness results it follows that the *control-to-state operator* \mathcal{S}

$$u \mapsto \mathcal{S}(u) := (\phi, \mu, \sigma)$$

is well-defined and Lipschitz continuous as a mapping from $\mathcal{U}_R \subset L^2(Q)$ into the following space

$$(L^\infty(0, T; (H^1(\Omega))' \cap L^2(0, T; H^1(\Omega))) \times L^2(0, T; (H^1(\Omega))') \times (L^\infty(0, T; (H^1(\Omega))' \cap L^2(Q))).$$

The triplet (ϕ, μ, σ) is the unique weak solution to the state system with data (ϕ_0, σ_0, u) over the time interval $[0, T]$. For convenience, we use the notations $\phi = \mathcal{S}_1(u)$ and $\sigma = \mathcal{S}_3(u)$ for the first and third component of $\mathcal{S}(u)$. Then we prove the following result that implies the existence of a solution to problem (CP).

Theorem (Existence of the optimal control)

Assume that (P1), (F1), (U1) and (C1)–(C3) are satisfied. Let $\phi_0 \in H_N^2(\Omega) \cap H^3(\Omega)$ and $\sigma_0 \in H^1(\Omega)$. Then there exists at least one minimizer $(\phi_*, \sigma_*, u_*, \tau_*)$ to problem (CP).

Namely, $\phi_* = \mathcal{S}_1(u_*)$, $\sigma_* = \mathcal{S}_3(u_*)$ satisfy

$$\mathcal{J}(\phi_*, \sigma_*, u_*, \tau_*) = \inf_{\substack{(w, s) \in \mathcal{U}_{\text{ad}} \times [0, T] \\ \text{s.t. } \phi = \mathcal{S}_1(w), \sigma = \mathcal{S}_3(w)}} \mathcal{J}(\phi, \sigma, w, s).$$

Differentiability of the control-to-state map

We establish then the Fréchet differentiability of the solution operator \mathcal{S} with respect to the control u .

Differentiability of the control-to-state map

We establish then the Fréchet differentiability of the solution operator \mathcal{S} with respect to the control u . For $u_* \in \mathcal{U}_R$, let $(\phi_*, \mu_*, \sigma_*) = \mathcal{S}(u_*)$. We consider for any $h \in L^2(Q)$ the linearized system

$$\begin{aligned}\partial_t \xi - \Delta \eta &= P'(\phi_*)(\sigma_* - \mu_*)\xi + P(\phi_*)(\rho - \eta), & \eta &= -\Delta \xi + F''(\phi_*)\xi, \\ \partial_t \rho - \Delta \rho &= -P'(\phi_*)(\sigma_* - \mu_*)\xi - P(\phi_*)(\rho - \eta) + h \\ \partial_n \xi &= \partial_n \eta = \partial_n \rho = 0, & \xi(0) &= \rho(0) = 0.\end{aligned}$$

Differentiability of the control-to-state map

We establish then the Fréchet differentiability of the solution operator \mathcal{S} with respect to the control u . For $u_* \in \mathcal{U}_R$, let $(\phi_*, \mu_*, \sigma_*) = \mathcal{S}(u_*)$. We consider for any $h \in L^2(Q)$ the linearized system

$$\begin{aligned}\partial_t \xi - \Delta \eta &= P'(\phi_*)(\sigma_* - \mu_*)\xi + P(\phi_*)(\rho - \eta), & \eta &= -\Delta \xi + F''(\phi_*)\xi, \\ \partial_t \rho - \Delta \rho &= -P'(\phi_*)(\sigma_* - \mu_*)\xi - P(\phi_*)(\rho - \eta) + h \\ \partial_n \xi &= \partial_n \eta = \partial_n \rho = 0, & \xi(0) &= \rho(0) = 0.\end{aligned}$$

We can apply [Theorems 3.1, 3.2, CGRS] for the well-posedness of the linearized system and the Fréchet differentiability of the control-to-state operator \mathcal{S} with respect to u .

Differentiability of the control-to-state map

We establish then the Fréchet differentiability of the solution operator \mathcal{S} with respect to the control u . For $u_* \in \mathcal{U}_R$, let $(\phi_*, \mu_*, \sigma_*) = \mathcal{S}(u_*)$. We consider for any $h \in L^2(Q)$ the linearized system

$$\begin{aligned}\partial_t \xi - \Delta \eta &= P'(\phi_*)(\sigma_* - \mu_*)\xi + P(\phi_*)(\rho - \eta), & \eta &= -\Delta \xi + F''(\phi_*)\xi, \\ \partial_t \rho - \Delta \rho &= -P'(\phi_*)(\sigma_* - \mu_*)\xi - P(\phi_*)(\rho - \eta) + h \\ \partial_n \xi &= \partial_n \eta = \partial_n \rho = 0, & \xi(0) &= \rho(0) = 0.\end{aligned}$$

We can apply [Theorems 3.1, 3.2, CGRS] for the well-posedness of the linearized system and the Fréchet differentiability of the control-to-state operator \mathcal{S} with respect to u . Assume (P1), (F1), (U1), (C1)–(C3), let $\phi_0 \in H_N^2(\Omega) \cap H^3(\Omega)$ and $\sigma_0 \in H^1(\Omega)$. Then the control-to-state operator \mathcal{S} is Fréchet differentiable in \mathcal{U}_R as a mapping from $L^2(Q)$ into

$$\begin{aligned}\mathcal{Y} &:= \left(H^1(0, T; (H_N^2(\Omega))') \cap L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H_N^2(\Omega)) \right) \times L^2(Q) \\ &\quad \times \left(H^1(0, T; L^2(\Omega)) \cap L^2(0, T; H^2(\Omega)) \right).\end{aligned}$$

Differentiability of the control-to-state map

We establish then the Fréchet differentiability of the solution operator \mathcal{S} with respect to the control u . For $u_* \in \mathcal{U}_R$, let $(\phi_*, \mu_*, \sigma_*) = \mathcal{S}(u_*)$. We consider for any $h \in L^2(Q)$ the linearized system

$$\begin{aligned}\partial_t \xi - \Delta \eta &= P'(\phi_*)(\sigma_* - \mu_*)\xi + P(\phi_*)(\rho - \eta), & \eta &= -\Delta \xi + F''(\phi_*)\xi, \\ \partial_t \rho - \Delta \rho &= -P'(\phi_*)(\sigma_* - \mu_*)\xi - P(\phi_*)(\rho - \eta) + h \\ \partial_n \xi &= \partial_n \eta = \partial_n \rho = 0, & \xi(0) &= \rho(0) = 0.\end{aligned}$$

We can apply [Theorems 3.1, 3.2, CGRS] for the well-posedness of the linearized system and the Fréchet differentiability of the control-to-state operator \mathcal{S} with respect to u . Assume (P1), (F1), (U1), (C1)–(C3), let $\phi_0 \in H_N^2(\Omega) \cap H^3(\Omega)$ and $\sigma_0 \in H^1(\Omega)$. Then the control-to-state operator \mathcal{S} is Fréchet differentiable in \mathcal{U}_R as a mapping from $L^2(Q)$ into

$$\begin{aligned}\mathcal{Y} &:= \left(H^1(0, T; (H_N^2(\Omega))') \cap L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H_N^2(\Omega)) \right) \times L^2(Q) \\ &\quad \times \left(H^1(0, T; L^2(\Omega)) \cap L^2(0, T; H^2(\Omega)) \right).\end{aligned}$$

For any $u_* \in \mathcal{U}_R$, the Fréchet derivative $D\mathcal{S}(u_*) \in \mathcal{L}(L^2(Q), \mathcal{Y})$ is defined as follows: for any $h \in L^2(Q)$, $D\mathcal{S}(u_*)h = (\xi^h, \eta^h, \rho^h)$, where (ξ^h, η^h, ρ^h) is the unique solution to the linearized system associated with h .

First order optimality conditions

Define a reduced functional

$$\tilde{\mathcal{J}}(u, \tau) := \mathcal{J}(\mathcal{S}_1(u), \mathcal{S}_3(u), u, \tau).$$

Since the control-to-state mapping \mathcal{S} is also Fréchet differentiable into $C^0([0, T]; L^2(\Omega))$ with respect to u , then the reduced cost functional $\tilde{\mathcal{J}}$ is Fréchet differentiable in \mathcal{U}_R .

First order optimality conditions

Define a reduced functional

$$\tilde{\mathcal{J}}(u, \tau) := \mathcal{J}(\mathcal{S}_1(u), \mathcal{S}_3(u), u, \tau).$$

Since the control-to-state mapping \mathcal{S} is also Fréchet differentiable into $C^0([0, T]; L^2(\Omega))$ with respect to u , then the reduced cost functional $\tilde{\mathcal{J}}$ is Fréchet differentiable in \mathcal{U}_R .

Theorem (Existence of solutions to the adjoint system)

Assume (P1), (F1), (U1), (C1)–(C3), $\phi_0 \in H_N^2(\Omega) \cap H^3(\Omega)$, and $\sigma_0 \in H^1(\Omega)$. Then the adjoint system

$$\begin{aligned} -\partial_t p + \Delta q - F''(\phi_*) q + P'(\phi_*)(\sigma_* - \mu_*)(r - p) &= \beta_Q (\phi_* - \phi_Q) \\ q - \Delta p + P(\phi_*)(p - r) = 0, \quad -\partial_t r - \Delta r + P(\phi_*)(r - p) &= \alpha_Q (\sigma_* - \sigma_Q) \\ \partial_n p = \partial_n q = \partial_n r = 0, \quad r(\tau_*) = 0, \quad p(\tau_*) &= \beta_\Omega (\phi_*(\tau_*) - \phi_\Omega) + \frac{\beta_S}{2} \end{aligned}$$

has a unique weak solution (p, q, r) on $[0, T]$:

$$\begin{aligned} p &\in H^1(0, T; (H_N^2(\Omega))') \cap C^0([0, T]; L^2(\Omega)) \cap L^2(0, T; H_N^2(\Omega)), \\ q &\in L^2(Q), \quad r \in H^1(0, T; L^2(\Omega)) \cap C^0([0, T]; H^1(\Omega)) \cap L^2(0, T; H_N^2(\Omega)). \end{aligned}$$

Necessary optimality conditions

Necessary optimality conditions

Theorem (Necessary optimality conditions)

Let $(u_*, \tau_*) \in \mathcal{U}_{\text{ad}} \times [0, T]$ denote a minimizer to the optimal control problem (CP) with corresponding state variables $(\phi_*, \mu_*, \sigma_*) = \mathcal{S}(u_*)$ and associated adjoint variables (p, q, r) , then it holds:

$$\beta_u \int_0^T \int_{\Omega} u_*(u - u_*) \, dx \, dt + \int_0^{\tau_*} \int_{\Omega} r(u - u_*) \, dx \, dt \geq 0, \quad \forall u \in \mathcal{U}_{\text{ad}}.$$

Besides, setting

$$\begin{aligned} \mathcal{L}(\phi_*, \sigma_*, \tau_*) &= \frac{\beta_Q}{2} \int_{\Omega} |\phi_*(\tau_*) - \phi_Q(\tau_*)|^2 \, dx + \beta_{\Omega} \int_{\Omega} (\phi_*(\tau_*) - \phi_{\Omega}) \partial_t \phi_*(\tau_*) \, dx \\ &+ \frac{\alpha_Q}{2} \int_{\Omega} |\sigma_*(\tau_*) - \sigma_Q|^2 \, dx + \frac{\beta_S}{2} \int_{\Omega} \partial_t \phi_*(\tau_*) \, dx + \beta_T \end{aligned}$$

we have

$$\mathcal{L}(\phi_*, \sigma_*, \tau_*) \begin{cases} \geq 0, & \text{if } \tau_* = 0, \\ = 0, & \text{if } \tau_* \in (0, T), \\ \leq 0, & \text{if } \tau_* = T. \end{cases}$$

Interpretation of the first condition

Besides, if we extend r by zero to $(\tau_*, T]$, then we can express the variational inequality

$$\beta_u \int_0^T \int_{\Omega} u_*(u - u_*) \, dx \, dt + \int_0^{\tau_*} \int_{\Omega} r(u - u_*) \, dx \, dt \geq 0, \quad \forall u \in \mathcal{U}_{\text{ad}}.$$

as

$$\int_0^T \int_{\Omega} (\beta_u u_* + r)(u - u_*) \, dx \, dt \geq 0, \quad \forall u \in \mathcal{U}_{\text{ad}},$$

which allows the interpretation that the optimal control u_* is the $L^2(Q)$ -projection of $-\beta_u^{-1}r$ onto the set \mathcal{U}_{ad} (provided that $\beta_u > 0$).

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna**
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Long-time behavior for a different model – joint work with A. Miranville
and G. Schimperna, arXiv:1810.12239v1 (2018)

Long-time behavior for a different model – joint work with A. Miranville and G. Schimperna, arXiv:1810.12239v1 (2018)

We consider here the long time dynamics for the following model for tumor growth:

$$\begin{aligned}\varphi_t - \Delta\mu &= (\mathcal{P}\sigma - \mathcal{A})h(\varphi), \\ \mu &= -\Delta\varphi + \Psi'(\varphi), \\ \sigma_t - \Delta\sigma &= -\mathcal{C}\sigma h(\varphi) + B(\sigma_s - \sigma),\end{aligned}$$

settled in $\Omega \times (0, +\infty)$, and complemented with the Cauchy conditions and with no-flux (i.e., homogeneous Neumann) boundary conditions for all unknowns.

Long-time behavior for a different model – joint work with A. Miranville and G. Schimperna, arXiv:1810.12239v1 (2018)

We consider here the long time dynamics for the following model for tumor growth:

$$\begin{aligned}\varphi_t - \Delta\mu &= (\mathcal{P}\sigma - \mathcal{A})h(\varphi), \\ \mu &= -\Delta\varphi + \Psi'(\varphi), \\ \sigma_t - \Delta\sigma &= -\mathcal{C}\sigma h(\varphi) + B(\sigma_s - \sigma),\end{aligned}$$

settled in $\Omega \times (0, +\infty)$, and complemented with the Cauchy conditions and with no-flux (i.e., homogeneous Neumann) boundary conditions for all unknowns.

- Here $h(s)$ is an interpolation function such that $h(-1) = 0$ and $h(1) = 1$, and
 - ▶ $h(\varphi)\mathcal{P}\sigma$ - proliferation of tumor cells proportional to nutrient concentration
 - ▶ $h(\varphi)\mathcal{A}$ - apoptosis of tumor cells
 - ▶ $h(\varphi)\mathcal{C}\sigma$ - consumption of nutrient by the tumor cells
- The constant σ_s denotes the nutrient concentration in a pre-existing vasculature, and $B(\sigma_s - \sigma)$ models the supply of nutrient from the blood vessels if $\sigma_s > \sigma$ and the transport of nutrient away from the domain Ω if $\sigma_s < \sigma$.

Long-time behavior for a different model – joint work with A. Miranville and G. Schimperna, arXiv:1810.12239v1 (2018)

We consider here the long time dynamics for the following model for tumor growth:

$$\begin{aligned}\varphi_t - \Delta\mu &= (\mathcal{P}\sigma - \mathcal{A})h(\varphi), \\ \mu &= -\Delta\varphi + \Psi'(\varphi), \\ \sigma_t - \Delta\sigma &= -\mathcal{C}\sigma h(\varphi) + B(\sigma_s - \sigma),\end{aligned}$$

settled in $\Omega \times (0, +\infty)$, and complemented with the Cauchy conditions and with no-flux (i.e., homogeneous Neumann) boundary conditions for all unknowns.

- Here $h(s)$ is an interpolation function such that $h(-1) = 0$ and $h(1) = 1$, and
 - ▶ $h(\varphi)\mathcal{P}\sigma$ - proliferation of tumor cells proportional to nutrient concentration
 - ▶ $h(\varphi)\mathcal{A}$ - apoptosis of tumor cells
 - ▶ $h(\varphi)\mathcal{C}\sigma$ - consumption of nutrient by the tumor cells
- The constant σ_s denotes the nutrient concentration in a pre-existing vasculature, and $B(\sigma_s - \sigma)$ models the supply of nutrient from the blood vessels if $\sigma_s > \sigma$ and the transport of nutrient away from the domain Ω if $\sigma_s < \sigma$.
- A regular double-well potential Ψ , e.g., $\Psi(s) = 1/4(1 - s^2)^2$

Long-time behavior for a different model – joint work with A. Miranville
and G. Schimperna, arXiv:1810.12239v1 (2018)

Long-time behavior for a different model – joint work with A. Miranville and G. Schimperna, arXiv:1810.12239v1 (2018)

The model was introduced in [Chen, Wise, Shenoy, Lowengrub (2014)] and the in [Garcke, Lam, Sitka, Styles (2016)] in a more general framework.

Long-time behavior for a different model – joint work with A. Miranville and G. Schimperna, arXiv:1810.12239v1 (2018)

The model was introduced in [Chen, Wise, Shenoy, Lowengrub (2014)] and then in [Garcke, Lam, Sitka, Styles (2016)] in a more general framework.

We prove that, under **physically motivated assumptions** on parameters and data,

- the corresponding initial-boundary value problem generates a dissipative dynamical system
- that admits the **global attractor** in a proper phase space.

Long-time behavior for a different model – joint work with A. Miranville and G. Schimperna, arXiv:1810.12239v1 (2018)

The model was introduced in [Chen, Wise, Shenoy, Lowengrub (2014)] and then in [Garcke, Lam, Sitka, Styles (2016)] in a more general framework.



We prove that, under **physically motivated assumptions** on parameters and data,

- the corresponding initial-boundary value problem generates a dissipative dynamical system
- that admits the **global attractor** in a proper phase space.

The main difference with respect to the previous model is that here we do not have the total energy balance we had before. Here we only have

$$\frac{d}{dt} \left(\frac{1}{2} \|\nabla \varphi\|^2 + \int_{\Omega} \Psi(\varphi) dx \right) + \|\nabla \mu\|^2 = \int_{\Omega} (\mathcal{P}\sigma - \mathcal{A})h(\varphi)\mu dx.$$

Examples of functions h and Ψ



The basic assumptions on the potential

The configuration potential Ψ lies in $C_{\text{loc}}^{1,1}(\mathbb{R})$. Moreover its derivative is decomposed as a sum of a **monotone part** β and a **linear perturbation**:

$$\Psi'(r) = \beta(r) - \lambda r, \quad \lambda \geq 0, \quad r \in \mathbb{R}.$$

We normalized so that $\beta(0) = 0$ and further β complies with the growth condition

$$\exists c_\beta > 0 : |\beta(r)| \leq c_\beta(1 + \Psi(r)) \quad \forall r \in \mathbb{R},$$

which is more or less equivalent to asking Ψ to have at most an exponential growth at infinity.

The basic assumptions on the potential

The configuration potential Ψ lies in $C_{\text{loc}}^{1,1}(\mathbb{R})$. Moreover its derivative is decomposed as a sum of a **monotone part** β and a **linear perturbation**:

$$\Psi'(r) = \beta(r) - \lambda r, \quad \lambda \geq 0, \quad r \in \mathbb{R}.$$

We normalized so that $\beta(0) = 0$ and further β complies with the growth condition

$$\exists c_\beta > 0 : |\beta(r)| \leq c_\beta(1 + \Psi(r)) \quad \forall r \in \mathbb{R},$$

which is more or less equivalent to asking Ψ to have at most an exponential growth at infinity. In order to avoid degenerate situations (such as $\beta = \Psi \equiv 0, \lambda = 0$) we also ask a minimal growth condition at infinity for Ψ , i.e. that

$$\liminf_{|r| \nearrow \infty} \frac{\Psi(r)}{|r|} =: \ell > 0.$$

The basic assumptions on the potential

The configuration potential Ψ lies in $C_{\text{loc}}^{1,1}(\mathbb{R})$. Moreover its derivative is decomposed as a sum of a **monotone part β** and a **linear perturbation**:

$$\Psi'(r) = \beta(r) - \lambda r, \quad \lambda \geq 0, \quad r \in \mathbb{R}.$$

We normalized so that $\beta(0) = 0$ and further β complies with the growth condition

$$\exists c_\beta > 0 : |\beta(r)| \leq c_\beta(1 + \Psi(r)) \quad \forall r \in \mathbb{R},$$

which is more or less equivalent to asking Ψ to have at most an exponential growth at infinity. In order to avoid degenerate situations (such as $\beta = \Psi \equiv 0$, $\lambda = 0$) we also ask a minimal growth condition at infinity for Ψ , i.e. that

$$\liminf_{|r| \nearrow \infty} \frac{\Psi(r)}{|r|} =: \ell > 0.$$

In order to prove uniqueness of solutions we also need that there exists $c > 0$ such that

$$|\beta(r) - \beta(s)| \leq c|r - s|(1 + |\beta(r)| + |\beta(s)|) \quad \forall r, s \in \mathbb{R}.$$

Note that this is still consistent with asking an **at most exponential growth of β** .

Assumptions on the function h and on the coefficients

The coefficients are assumed to satisfy $\mathcal{P}, \mathcal{A}, B, \mathcal{C} > 0$, $\sigma_c \in (0, 1)$.

Assumptions on the function h and on the coefficients

The coefficients are assumed to satisfy $\mathcal{P}, \mathcal{A}, B, \mathcal{C} > 0$, $\sigma_c \in (0, 1)$.

Next, we assume that

- h is in $C^1(\mathbb{R})$, increasingly monotone and it satisfies at least $h(-1) = 0$ and $h(r) \equiv 1$ for all $r \geq 1 \implies h$ is globally Lipschitz continuous

Assumptions on the function h and on the coefficients

The coefficients are assumed to satisfy $\mathcal{P}, \mathcal{A}, B, \mathcal{C} > 0$, $\sigma_c \in (0, 1)$.

Next, we assume that

- h is in $C^1(\mathbb{R})$, increasingly monotone and it satisfies at least $h(-1) = 0$ and $h(r) \equiv 1$ for all $r \geq 1 \implies h$ is globally Lipschitz continuous
- There exist $\underline{h} \geq 0$ and $\underline{\varphi} \leq -1$ such that $h(r) \equiv -\underline{h}$ for all $r \leq \underline{\varphi}$

Assumptions on the function h and on the coefficients

The coefficients are assumed to satisfy $\mathcal{P}, \mathcal{A}, B, \mathcal{C} > 0$, $\sigma_c \in (0, 1)$.

Next, we assume that

- h is in $C^1(\mathbb{R})$, increasingly monotone and it satisfies at least $h(-1) = 0$ and $h(r) \equiv 1$ for all $r \geq 1 \implies h$ is globally Lipschitz continuous
- There exist $\underline{h} \geq 0$ and $\underline{\varphi} \leq -1$ such that $h(r) \equiv -\underline{h}$ for all $r \leq \underline{\varphi}$

Remark

*The function $h(\varphi)$ is assumed to satisfy $h(-1) = 0$ and $h(1) = 1$. The simplest situation when this occurs is the “symmetric” case when we have $\underline{h} = 0$ and $\underline{\varphi} = -1$. On the other hand we will see in what follows that **dissipativity** of trajectories may not hold in such a case. This motivates our choice to consider the possibility of having $\underline{h} > 0$.*

Assumptions on the function h and on the coefficients

The coefficients are assumed to satisfy $\mathcal{P}, \mathcal{A}, B, \mathcal{C} > 0$, $\sigma_c \in (0, 1)$.

Next, we assume that

- h is in $C^1(\mathbb{R})$, increasingly monotone and it satisfies at least $h(-1) = 0$ and $h(r) \equiv 1$ for all $r \geq 1 \implies h$ is globally Lipschitz continuous
- There exist $\underline{h} \geq 0$ and $\underline{\varphi} \leq -1$ such that $h(r) \equiv -\underline{h}$ for all $r \leq \underline{\varphi}$

Remark

The function $h(\varphi)$ is assumed to satisfy $h(-1) = 0$ and $h(1) = 1$. The simplest situation when this occurs is the “symmetric” case when we have $\underline{h} = 0$ and $\underline{\varphi} = -1$. On the other hand we will see in what follows that **dissipativity** of trajectories may not hold in such a case. This motivates our choice to consider the possibility of having $\underline{h} > 0$.

Remark

We could also take $h(\varphi) = k\varphi + h_0(\varphi)$, where $k > 0$ and h_0 is smooth and uniformly bounded. This situation is somehow simpler because, at least as long as we can guarantee that $\mathcal{P}\sigma - \mathcal{A} > 0$, the linear part of h drives some mass dissipation effect in the Cahn-Hilliard type equation $\varphi_t - \Delta\mu = (\mathcal{P}\sigma - \mathcal{A})h(\varphi)$.

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness**
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

We assume the initial data to satisfy

$$\begin{aligned}\sigma_0 &\in L^\infty(\Omega), & 0 \leq \sigma_0 \leq 1 \text{ a.e. in } \Omega, \\ \varphi_0 &\in H^1(\Omega), & \Psi(\varphi_0) \in L^1(\Omega).\end{aligned}$$

We assume the initial data to satisfy

$$\begin{aligned}\sigma_0 &\in L^\infty(\Omega), & 0 \leq \sigma_0 \leq 1 \text{ a.e. in } \Omega, \\ \varphi_0 &\in H^1(\Omega), & \Psi(\varphi_0) \in L^1(\Omega).\end{aligned}$$

Theorem (Well-posedness)

Then the tumor-growth model

$$\begin{aligned}\varphi_t - \Delta \mu &= (\mathcal{P}\sigma - \mathcal{A})h(\varphi), & \varphi(0) &= \varphi_0, & \partial_n \varphi &= 0 \text{ on } \partial\Omega, \\ \mu &= -\Delta \varphi + \Psi'(\varphi), & \partial_n \mu &= 0 \text{ on } \partial\Omega, \\ \sigma_t - \Delta \sigma &= -C\sigma h(\varphi) + B(\sigma_s - \sigma), & \sigma(0) &= \sigma_0, & \partial_n \sigma &= 0 \text{ on } \partial\Omega\end{aligned}$$

admits *one and only one global in time weak solution*:

$$\begin{aligned}\varphi &\in H^1(0, T; H^1(\Omega)') \cap C^0([0, T]; H^1(\Omega)) \cap L^2(0, T; H^2(\Omega)), \\ \beta(\varphi) &\in L^2(0, T; L^2(\Omega)), \quad \mu \in L^2(0, T; H^1(\Omega)), \\ \sigma &\in H^1(0, T; H^1(\Omega)') \cap C^0([0, T]; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega)) \cap L^\infty(0, T; L^\infty(\Omega));\end{aligned}$$

Moreover, for any $T > 0$ there exists $\bar{\sigma}_T \geq 1$ such that

$$0 \leq \sigma(t, x) \leq \bar{\sigma}_T, \quad \text{for a.e. } (t, x) \in (0, T) \times \Omega,$$

where we can take $\bar{\sigma}_T$ independent of time if $B - Ch > 0$ and $\bar{\sigma}_T = 1$ if $h = 0$.

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor**
- 9 Perspectives and Open problems

Assumptions for dissipativity

Let the parameters in

$$\begin{aligned}\varphi_t - \Delta\mu &= (\mathcal{P}\sigma - \mathcal{A})h(\varphi), \\ \mu &= -\Delta\varphi + \Psi'(\varphi), \\ \sigma_t - \Delta\sigma &= -\mathcal{C}\sigma h(\varphi) + B(\sigma_s - \sigma),\end{aligned}$$

satisfy (where $h(r) \equiv -\underline{h}$ for all $r \leq \varphi \leq -1$)

$$(H1) \quad \underline{h} > 0, \quad B - \mathcal{C}\underline{h} > 0,$$

$$(H2) \quad \frac{B\sigma_s}{B - \mathcal{C}\underline{h}} < 1,$$

$$(H3) \quad \mathcal{A} - \mathcal{P} \frac{B\sigma_s}{B - \mathcal{C}\underline{h}} > 0.$$

Assumptions for dissipativity

Let the parameters in

$$\begin{aligned}\varphi_t - \Delta\mu &= (\mathcal{P}\sigma - \mathcal{A})h(\varphi), \\ \mu &= -\Delta\varphi + \Psi'(\varphi), \\ \sigma_t - \Delta\sigma &= -\mathcal{C}\sigma h(\varphi) + B(\sigma_s - \sigma),\end{aligned}$$

satisfy (where $h(r) \equiv -\underline{h}$ for all $r \leq \varphi \leq -1$)

$$(H1) \quad \underline{h} > 0, \quad B - \mathcal{C}\underline{h} > 0,$$

$$(H2) \quad \frac{B\sigma_s}{B - \mathcal{C}\underline{h}} < 1,$$

$$(H3) \quad \mathcal{A} - \mathcal{P} \frac{B\sigma_s}{B - \mathcal{C}\underline{h}} > 0.$$

These conditions essentially prescribe \underline{h} to be *strictly positive, but small*.

Assumptions for dissipativity

Let the parameters in

$$\begin{aligned}\varphi_t - \Delta\mu &= (\mathcal{P}\sigma - \mathcal{A})h(\varphi), \\ \mu &= -\Delta\varphi + \Psi'(\varphi), \\ \sigma_t - \Delta\sigma &= -\mathcal{C}\sigma h(\varphi) + B(\sigma_s - \sigma),\end{aligned}$$

satisfy (where $h(r) \equiv -\underline{h}$ for all $r \leq \varphi \leq -1$)

$$(H1) \quad \underline{h} > 0, \quad B - \mathcal{C}\underline{h} > 0,$$

$$(H2) \quad \frac{B\sigma_s}{B - \mathcal{C}\underline{h}} < 1,$$

$$(H3) \quad \mathcal{A} - \mathcal{P} \frac{B\sigma_s}{B - \mathcal{C}\underline{h}} > 0.$$

These conditions essentially prescribe \underline{h} to be *strictly positive, but small*.

Let also β have a superquadratic behavior at infinity, namely

$$\exists \kappa_\beta > 0, C_\beta \geq 0, p_\beta > 2 : \beta(r) \operatorname{sign} r \geq \kappa_\beta |r|^{p_\beta} - C_\beta \quad \forall r \in \mathbb{R}.$$

The spatially homogeneous case: (H1)

Starting from spatially homogeneous initial data we reduce to the following ODE system:

$$\begin{aligned}X' + (\mathcal{A} - \mathcal{P}S)h(X) &= 0, \\S' + \mathcal{C}Sh(X) + B(S - \sigma_s) &= 0\end{aligned}$$

where $X = X(t)$ and $S = S(t)$ are the spatial mean values of φ and σ .

The spatially homogeneous case: (H1)

Starting from spatially homogeneous initial data we reduce to the following ODE system:

$$\begin{aligned}X' + (\mathcal{A} - \mathcal{P}S)h(X) &= 0, \\S' + \mathcal{C}Sh(X) + B(S - \sigma_s) &= 0\end{aligned}$$

where $X = X(t)$ and $S = S(t)$ are the spatial mean values of φ and σ .

1) If $\underline{h} = 0$, i.e. **H1 i) does not hold**, and $X(0) < -1$ then $X(t)$ is conserved in time.

There is no hope to prove that $X(t)$ eventually lies in some bounded absorbing set.

The spatially homogeneous case: (H1)

Starting from spatially homogeneous initial data we reduce to the following ODE system:

$$\begin{aligned}X' + (\mathcal{A} - \mathcal{P}S)h(X) &= 0, \\S' + \mathcal{C}Sh(X) + B(S - \sigma_s) &= 0\end{aligned}$$

where $X = X(t)$ and $S = S(t)$ are the spatial mean values of φ and σ .

- 1) If $\underline{h} = 0$, i.e. **H1) i) does not hold**, and $X(0) < -1$ then $X(t)$ is conserved in time. There is no hope to prove that $X(t)$ eventually lies in some bounded absorbing set.
- 2) Let us now assume $\underline{h} > 0$. Then we have

$$B\sigma_s - (C + B)S \leq S' \leq B\sigma_s - (B - C\underline{h})S$$

The spatially homogeneous case: (H1)

Starting from spatially homogeneous initial data we reduce to the following ODE system:

$$\begin{aligned}X' + (\mathcal{A} - \mathcal{P}S)h(X) &= 0, \\S' + \mathcal{C}Sh(X) + B(S - \sigma_s) &= 0\end{aligned}$$

where $X = X(t)$ and $S = S(t)$ are the spatial mean values of φ and σ .

- 1) If $\underline{h} = 0$, i.e. **(H1) i) does not hold**, and $X(0) < -1$ then $X(t)$ is conserved in time. There is no hope to prove that $X(t)$ eventually lies in some bounded absorbing set.
- 2) Let us now assume $\underline{h} > 0$. Then we have

$$B\sigma_s - (C + B)S \leq S' \leq B\sigma_s - (B - C\underline{h})S$$

If $C\underline{h} \geq B$, i.e. **(H1) ii) does not hold** and $X(0) \ll 0$, $S(0) \gg 0$ (in such a way that $\mathcal{P}S - \mathcal{A} > 0$), then it follows

$$\begin{aligned}X' &= -(\mathcal{P}S - \mathcal{A})\underline{h} < 0, \\S' &= B\sigma_s + (C\underline{h} - B)S > 0\end{aligned}$$


and both $|X|$ and S go increasing forever. Even if we restrict ourselves to $S(0) \leq 1$, if $X(0) < -1$ then the physical constraint $S(t) \in [0, 1]$ is not respected.

The spatially homogeneous case: (H2)

Assume (H1): $\underline{h} > 0$, $B - C\underline{h} > 0$. Then, the region

$S := \left\{ (X, S) : \frac{B\sigma_s}{C+B} \leq S \leq \frac{B\sigma_s}{B-C\underline{h}} \right\}$ is positively invariant for the dynamical process

because $B\sigma_s - (C+B)S \leq S' \leq B\sigma_s - (B-C\underline{h})S$:




The spatially homogeneous case: (H2)

Assume (H1): $\underline{h} > 0$, $B - C\underline{h} > 0$. Then, the region

$S := \left\{ (X, S) : \frac{B\sigma_s}{C+B} \leq S \leq \frac{B\sigma_s}{B-C\underline{h}} \right\}$ is positively invariant for the dynamical process

because $B\sigma_s - (C+B)S \leq S' \leq B\sigma_s - (B-C\underline{h})S$:



Now, if we want to keep the physical constraint $S(t) \in [0, 1]$, we need to assume


$$(H2) : \frac{B\sigma_s}{B - C\underline{h}} < 1$$

The spatially homogeneous case: (H3)

Let us assume that $X(0) > 1$, which also implies $h(X) = 1$. Then, we have:

$$X' = (PS - A)$$

and condition (H3): $\frac{A}{P} > \frac{B\sigma_s}{B-Ch}$ prescribes that in $S \cap \{X > 1\}$ we have $X' < 0$:




The spatially homogeneous case: (H3)

Let us assume that $X(0) > 1$, which also implies $h(X) = 1$. Then, we have:

$$X' = (PS - A)$$

and condition (H3): $\frac{A}{P} > \frac{B\sigma_s}{B-Ch}$ prescribes that in $S \cap \{X > 1\}$ we have $X' < 0$:



On the other hand when $\frac{A}{P} \leq \frac{B\sigma_s}{B+C}$, dissipativity cannot hold. Indeed if

$S(0) \in \left[\frac{B\sigma_s}{C+B}, \frac{B\sigma_s}{B-Ch} \right]$ and $X(0) \geq 1$, then $X(t)$ is forced to increase forever ($X' > 0$).

Dissipativity and Attractor

We can define the “energy space”

$$\mathcal{X} := \{(\varphi, \sigma) \in H^1(\Omega) \times L^\infty(\Omega) : \Psi(\varphi) \in L^1(\Omega)\}$$

and we correspondingly introduce the “magnitude” of an element $(\varphi, \sigma) \in \mathcal{X}$ as

$$\|(\varphi, \sigma)\|_{\mathcal{X}} := \|\varphi\|_{H^1} + \|\sigma\|_{L^\infty} + \|\Psi(\varphi)\|_{L^1}.$$

Dissipativity and Attractor

We can define the “energy space”

$$\mathcal{X} := \{(\varphi, \sigma) \in H^1(\Omega) \times L^\infty(\Omega) : \Psi(\varphi) \in L^1(\Omega)\}$$

and we correspondingly introduce the “magnitude” of an element $(\varphi, \sigma) \in \mathcal{X}$ as

$$\|(\varphi, \sigma)\|_{\mathcal{X}} := \|\varphi\|_{H^1} + \|\sigma\|_{L^\infty} + \|\Psi(\varphi)\|_{L^1}.$$

Theorem (Dissipativity)

Under the previous compatibility conditions, there exists a positive constant C_0 independent of the initial data and a time T_0 depending only on the \mathcal{X} -magnitude of the initial data such that any weak solution satisfies

$$\|(\varphi(t), \sigma(t))\|_{\mathcal{X}} \leq C_0 \quad \text{for every } t \geq T_0.$$

Dissipativity and Attractor

We can define the “energy space”

$$\mathcal{X} := \{(\varphi, \sigma) \in H^1(\Omega) \times L^\infty(\Omega) : \Psi(\varphi) \in L^1(\Omega)\}$$

and we correspondingly introduce the “magnitude” of an element $(\varphi, \sigma) \in \mathcal{X}$ as

$$\|(\varphi, \sigma)\|_{\mathcal{X}} := \|\varphi\|_{H^1} + \|\sigma\|_{L^\infty} + \|\Psi(\varphi)\|_{L^1}.$$

Theorem (Dissipativity)

Under the previous compatibility conditions, there exists a positive constant C_0 independent of the initial data and a time T_0 depending only on the \mathcal{X} -magnitude of the initial data such that any weak solution satisfies

$$\|(\varphi(t), \sigma(t))\|_{\mathcal{X}} \leq C_0 \quad \text{for every } t \geq T_0.$$

Theorem (Existence of the Attractor)

*Under the previous compatibility conditions the dynamical system generated by weak trajectories on the phase space \mathcal{X} admits the **global attractor** \mathcal{A} . More precisely, \mathcal{A} is a relatively compact subset of \mathcal{X} which is also bounded in $H^2(\Omega) \times H^1(\Omega)$ and uniformly attracts the trajectories emanating from any bounded set $\mathcal{B} \subset \mathcal{X}$.*

Outline

- 1 Phase field models for tumor growth
- 2 Recent joint work with C. Cavaterra and H. Wu
- 3 Well-posedness
- 4 Long-term dynamics
- 5 The optimal control problem
- 6 Recent joint work with A. Miranville and G. Schimperna
- 7 Well-posedness
- 8 Dissipativity and existence of the attractor
- 9 Perspectives and Open problems

Perspectives and Open problems

1. To study the **long-time behavior of solutions in terms of attractors**: with A. Miranville and G. Schimperna (on a further generalization of the model proposed by H. Garcke et. al. including **chemotaxis**), with A. Giorgini, K.-F. Lam, and G. Schimperna (attractors for a model including **velocities** proposed by Lowengrub et al.).

Perspectives and Open problems

1. To study the **long-time behavior of solutions in terms of attractors**: with A. Miranville and G. Schimperna (on a further generalization of the model proposed by H. Garcke et. al. including **chemotaxis**), with A. Giorgini, K.-F. Lam, and G. Schimperna (attractors for a model including **velocities** proposed by Lowengrub et al.).
2. **The study of optimal control: for a prostate model** introduced by H. Gomez et al. and proposed to us by G. Lorenzo and A. Reali (ongoing project with P. Colli and G. Marinoschi).

Perspectives and Open problems

1. To study the **long-time behavior of solutions in terms of attractors**: with A. Miranville and G. Schimperna (on a further generalization of the model proposed by H. Garcke et. al. including **chemotaxis**), with A. Giorgini, K.-F. Lam, and G. Schimperna (attractors for a model including **velocities** proposed by Lowengrub et al.).
2. **The study of optimal control: for a prostate model** introduced by H. Gomez et al. and proposed to us by G. Lorenzo and A. Reali (ongoing project with P. Colli and G. Marinoschi).
3. **To add the mechanics in Lagrangean coordinates** in a multiphase model: for example considering the tumor sample as a **porous media** (ongoing project with P. Krejčí and J. Sprekels).

Perspectives and Open problems

1. To study the **long-time behavior of solutions in terms of attractors**: with A. Miranville and G. Schimperna (on a further generalization of the model proposed by H. Garcke et. al. including **chemotaxis**), with A. Giorgini, K.-F. Lam, and G. Schimperna (attractors for a model including **velocities** proposed by Lowengrub et al.).
2. **The study of optimal control: for a prostate model** introduced by H. Gomez et al. and proposed to us by G. Lorenzo and A. Reali (ongoing project with P. Colli and G. Marinoschi).
3. **To add the mechanics in Lagrangean coordinates** in a multiphase model: for example considering the tumor sample as a **porous media** (ongoing project with P. Krejčí and J. Sprekels).
4. **Include a stochastic** term in phase-field models for tumor growth representing for example uncertainty of a therapy or random oscillations of the tumor phase (ongoing project with C. Orrieri and L. Scarpa).

Many thanks to all of you for the attention!

- Def. \mathcal{B}_0 is an *absorbing set* for a semigroup $S(t)$ on a metric space (X, d_X) iff
 - ▶ \mathcal{B}_0 is bdd
 - ▶ $\forall B \subset X$ bdd $\exists T_B \geq 0$ s.t. $S(t)B \subset \mathcal{B}_0 \quad \forall t \geq T_B$.
- Theorem. Let $S(t)$ be a strongly continuous semigroup on a c.m.s. (X, d_X) .
Moreover, if
 - ▶ $S(t)$ admits an absorbing set \mathcal{B}_0 ;
 - ▶ $\forall B \subset X$ bdd $\exists t_B > 0$ s.t. $\bigcup_{t \geq t_B} S(t)B$ is compact in X ,then $S(t)$ admits a *universal attractor* \mathcal{A} that is

$$\mathcal{A} = \bigcap_{\tau \geq 0} \overline{\bigcup_{t \geq \tau} S(t)\mathcal{B}_0}.$$